Ammoperdix griseogularis (See-see Partridge)

European Red List of Birds

Supplementary Material

The European Union (EU28) Red List assessments were based principally on the official data reported by EU Member States to the European Commission under Article 12 of the Birds Directive in 2019-20. For the European Red List assessments, similar data were sourced from BirdLife Partners and other collaborating experts in other European countries and territories. For more information, see BirdLife International (2021).

Contents

Reported national population sizes and trends
Trend maps of reported national population data
Sources of reported national population data
Species factsheet bibliography

Recommended citation

BirdLife International (2021) European Red List of Birds. Luxembourg: Publications Office of the European Union.

Further information

http://datazone.birdlife.org/info/euroredlist
http://www.birdlife.org/europe-and-central-asia/european-red-list-birds-0
http://www.iucnredlist.org/regions/europe
http://ec.europa.eu/environment/nature/conservation/species/redlist/

Data requests and feedback

To request access to these data in electronic format, provide new information, correct any errors or provide feedback, please email science@birdlife.org.

Ammoperdix griseogularis (See-see Partridge)

Table 1. Reported national breeding population size and trends in Europe[1].

Country (or territory)[2]	Population estimate				Short-term population trend[5]				Long-term population trend[5]				Subspecific population (where relevant)
	Size (pairs)[3]	Europe (%)	Year(s)	Method[4]	Direction[6]	Magnitude (%)[7]	Year(s)	Method[4]	Direction[6]	Magnitude (%)[7]	Year(s)	Method[4]	
Armenia	5–10	<1	2013-2018	complete	F		2007-2018	expert	?		2003-2018	deficient	
Azerbaijan	200–1000	9	1996-2019	expert	+		2013-2019	expert	+		1980-2019	complete	
Turkey	3700–5800	91	2013-2019	expert	-		2012-2018	expert	?		2010-2019	expert	
Europe	**3900–6800**	**100**											

[1] See 'Sources' at end of factsheet, and for more details on individual EU Member State reports, see the Article 12 reporting portal at http://bd.eionet.europa.eu/article12/report.
[2] The designation of geographical entities and the presentation of the material do not imply the expression of any opinion whatsoever on the part of IUCN or BirdLife International concerning the legal status of any country, territory or area, or of its authorities, or concerning the delimitation of its frontiers or boundaries.
[3] In the few cases where population size estimates were reported in units other than those specified, they were converted to the correct units using standard correction factors.
[4] The 'method used' (replacing the data 'quality' assessment in the 2015 European Red List) is reported as: a) Complete: complete survey or a statistically robust estimate; b) Partial: based mainly on extrapolation from a limited amount of data; c) Expert: based mainly on expert opinion with very limited data; d) Defficient: insufficient or no data available.
[5] The robustness of regional trends to the effects of any missing or incomplete data was tested using plausible scenarios, based on other sources of information, including any other reported information, recent national Red Lists, scientific literature, other publications and consultation with relevant experts.
[6] Trend directions are reported as: increasing (+); decreasing (-); stable (0); fluctuating (F); or unknown (?).
[7] Trend magnitudes are rounded to the nearest integer.

Trend maps

A symbol appears in each country where the species occurs: the shape and colour of the symbol represent the population trend in that country, and the size of the symbol corresponds to the proportion of the European population occurring in that country.

KEY

- Large increase (≥50%)
- Moderate increase (20–49%)
- Small increase (<20%)
- Increase of unknown magnitude
- Large decrease (≥50%)
- Moderate decrease (20–49%)
- Small decrease (<20%)
- Decrease of unknown magnitude
- Stable or fluctuating
- Unknown
- Present (no population or trend data)
- Extinct since 1980

Each symbol, with the exception of Present and Extinct, may occur in up to three different size classes, corresponding to the proportion of the European population occurring in that country.

- Large: ≥10% of the European population
- Medium: 1–9% of the European population
- Small: <1% of the European population

The designation of geographical entities and the presentation of the material do not imply the expression of any opinion whatsoever on the part of IUCN or BirdLife International concerning the legal status of any country, territory or area, or of its authorities, or concerning the delimitation of its frontiers or boundaries.

Figure 1. Breeding population sizes and short-term trends across Europe.



Figure 2. Breeding population sizes and long-term trends across Europe.



Sources

Armenia

Breeding population size: TSE NGO National Bird Monitoring data.
Breeding short-term trend: TSE (2020) The Atlas of the Breeding Birds in Armenia. In preparation.
Breeding long-term trend: TSE (2020) The Atlas of the Breeding Birds in Armenia. In preparation.

Azerbaijan

Breeding population size: AOS Data Base
Breeding short-term trend: AOS data base
Breeding long-term trend: AOS Data Base

Turkey

Breeding population size: Emin Yoğutçuoğlu and Levent Erkol personal communication (2019),, Kusbank Bird Database (Ebird) Kirwan G.M., Boyla K. A., Castell P., Demirci B., Özen M., Welch H., Marlow T., 2008, Birds of Turkey. Londra, Christopher Helm, 978-1-4081-0475-3,32
Breeding short-term trend: Ferdi Akarsu Pers. Communication and Birdlife Turkey Ceylanpınar, Akçakale and Kızılkuyu Semidesert areas reports.
Breeding long-term trend: Ferdi Akarsu Pers. Communication and Birdlife Turkey Ceylanpınar, Akçakale and Kızılkuyu Semidesert areas reports.

Bibliography

Bird, J. P., Martin, R., Akçakaya, H. R., Gilroy, J., Burfield, I. J., Garnett, S. G., Symes, A., Taylor, J., Sekercioglu, Ç. H. and Butchart, S. H. M. 2020. Generation lengths of the world's birds and their implications for extinction risk. Conservation Biology 34(5): 1252-1261. DOI: 10.1111/cobi.13486.

BirdLife International. 2004. Birds in Europe: population estimates, trends and conservation status. BirdLife International, Cambridge, U.K.

Biricik, M. and Karakas, R. 2012. Birds of Hasankeyf (South-Eastern Anatolia, Turkey) Under the Threat of a Big Dam Project. Natural Areas Journal 32(1): 96-105.

Fuller, R.A., Carroll, J.P. and McGowan, P.J.K. 2000. Partridges, Quails, Francolins, Snowcocks, Guineafowl, and Turkeys. Status Survey and Conservation Action Plan 2000 –2004. IUCN and World Pheasant Association, Gland, Switzerland and Cambridge, UK.

Keller, V., Herrando, S., Voříšek, P., Franch, M., Kipson, M., Milanesi, P., Martí, D., Anton, M., Klvanová, A., Kalyakin, M.V., Bauer, H.-G. and Foppen, R.P.B. 2020. European Breeding Bird Atlas 2: Distribution, Abundance and Change. European Bird Census Council & Lynx Edicions, Barcelona.

Madge, S. and McGowan, P. 2002. Pheasants, partridges and grouse: Including buttonquails, sandgrouse and allies. Christopher Helm, London.

McGowan, P.J.K. and Kirwan, G.M. 2014. See-see Partridge (Ammoperdix griseogularis). In: J. del Hoyo, A. Elliott, J. Sargatal, D.A. Christie and E. de Juana (eds), Handbook of the Birds of the World Alive, Lynx Edicions, Barcelona.

Norouzi, M., Mansouri, B., Hamidian, A.H., Ebrahimi, T. and Kardoni, F. 2012. Comparison of the metal concentrations in the feathers of three bird species from Southern Iran. Bulletin of environmental contamination and toxicology 89(5): 1082-1086.

Zolfaghari, G., Esmaili-Sari, A., Ghasempouri, S.M. and Kiabi, B.H. 2007. Examination of mercury concentration in the feathers of 18 species of birds in southwest Iran. Environmental Research 104(2): 258-265.