

***Glareola nordmanni* (Black-winged Pratincole)**

European Red List of Birds

Supplementary Material

The European Union (EU28) Red List assessments were based principally on the official data reported by EU Member States to the European Commission under Article 12 of the Birds Directive in 2019-20. For the European Red List assessments, similar data were sourced from BirdLife Partners and other collaborating experts in other European countries and territories. For more information, see BirdLife International (2021).

Contents

Reported national population sizes and trends
Trend maps of reported national population data
Sources of reported national population data
Species factsheet bibliography

Recommended citation

BirdLife International (2021) European Red List of Birds. Luxembourg: Publications Office of the European Union.

Further information

<http://datazone.birdlife.org/info/euroredlist>
<http://www.birdlife.org/europe-and-central-asia/european-red-list-birds-0>
<http://www.iucnredlist.org/regions/europe>
<http://ec.europa.eu/environment/nature/conservation/species/redlist/>

Data requests and feedback

To request access to these data in electronic format, provide new information, correct any errors or provide feedback, please email science@birdlife.org.

Glareola nordmanni (Black-winged Pratincole)

Table 1. Reported national breeding population size and trends in Europe¹.

Country (or territory) ²	Population estimate				Short-term population trend ⁵				Long-term population trend ⁵				Subspecific population (where relevant)
	Size (pairs) ³	Europe (%)	Year(s)	Method ⁴	Direction ⁶	Magnitude (%) ⁷	Year(s)	Method ⁴	Direction ⁶	Magnitude (%) ⁷	Year(s)	Method ⁴	
Russia	6000–12000	100	2008-2018	partial	0		2008-2018	expert	0		1980-2018	expert	
Turkey	0–49	<1	2002-2012	expert	?		2008-2019	deficient	?		1980-2013	deficient	
Ukraine	10	<1	2013-2018	expert	-		2009-2018	expert	-		1994-2018	expert	
EU28	0	<1											
Europe	6000–12100	100											

¹ See 'Sources' at end of factsheet, and for more details on individual EU Member State reports, see the Article 12 reporting portal at <http://bd.eionet.europa.eu/article12/report>.

² The designation of geographical entities and the presentation of the material do not imply the expression of any opinion whatsoever on the part of IUCN or BirdLife International concerning the legal status of any country, territory or area, or of its authorities, or concerning the delimitation of its frontiers or boundaries.

³ In the few cases where population size estimates were reported in units other than those specified, they were converted to the correct units using standard correction factors.

⁴ The 'method used' (replacing the data 'quality' assessment in the 2015 European Red List) is reported as: a) Complete: complete survey or a statistically robust estimate; b) Partial: based mainly on extrapolation from a limited amount of data; c) Expert: based mainly on expert opinion with very limited data; d) Defficient: insufficient or no data available.

⁵ The robustness of regional trends to the effects of any missing or incomplete data was tested using plausible scenarios, based on other sources of information, including any other reported information, recent national Red Lists, scientific literature, other publications and consultation with relevant experts.

⁶ Trend directions are reported as: increasing (+); decreasing (-); stable (0); fluctuating (F); or unknown (?).

⁷ Trend magnitudes are rounded to the nearest integer.

Trend maps

A symbol appears in each country where the species occurs: the shape and colour of the symbol represent the population trend in that country, and the size of the symbol corresponds to the proportion of the European population occurring in that country.

KEY

- | | |
|---|----------------------------------|
| ↑ Large increase ($\geq 50\%$) | ↓ Large decrease ($\geq 50\%$) |
| ↑ Moderate increase (20–49%) | ↓ Moderate decrease (20–49%) |
| ↑ Small increase ($< 20\%$) | ↓ Small decrease ($< 20\%$) |
| ↑ Increase of unknown magnitude | ↓ Decrease of unknown magnitude |
| ■ Stable or fluctuating | |
| □ Unknown | |
| ○ Present (no population or trend data) | |
| × Extinct since 1980 | |

Each symbol, with the exception of Present and Extinct, may occur in up to three different size classes, corresponding to the proportion of the European population occurring in that country.

- ↑ Large: $\geq 10\%$ of the European population
- ↑ Medium: 1–9% of the European population
- ↑ Small: $< 1\%$ of the European population

The designation of geographical entities and the presentation of the material do not imply the expression of any opinion whatsoever on the part of IUCN or BirdLife International concerning the legal status of any country, territory or area, or of its authorities, or concerning the delimitation of its frontiers or boundaries.

Figure 1. Breeding population sizes and short-term trends across Europe.





Figure 2. Breeding population sizes and long-term trends across Europe.



Sources

Russia

Breeding population size: Belik 2014; in prep; Database of the project on Atlas of breeding birds of European Russia
Breeding short-term trend: Fedosov & Malovichko 2006; Malovichko & Fedosov 2008; Belik in prep.
Breeding long-term trend: Belik et al. 2003; Belik 2004a,b; Malovichko & Fedosov 2008

Turkey

Breeding population size: Soner Bekir, Murat Bozdoğan personal communication (2019), Kirwan G.M., Boyla K. A., Castell P., Demirci B., Özen M., Welch H., Marlow T., 2008, Birds of Turkey. Londra, Christopher Helm, 978-1-4081-0475- Eken G., Bozdoğan M., İsfendiyaroğlu S., Kılıç D.T., Lise Y. (2006) Türkiye'nin Önemli Doğa Alanları. Doğa Derneği, Ankara. Kılıç DT, Eken G. 2004. Türkiye'nin önemli kus alanları—2004 güncellemesi (Important bird areas in Turkey—2004 update). Ankara: Doga Derneği.

Ukraine

Breeding population size: Atlas work, non-published data
Breeding short-term trend: КіндаВ.В., 2009, Дерихвіст степовий. Червона книга України. Тваринний світ. К., С. 458.
Breeding long-term trend: Горбань І. (2003): Оцінка чисельності гніздових птахів України. - Вісн. Львів. ун-ту. Сер. біол. 34: 147-158.

Bibliography

- Ash, J. S. 1977. Four species of birds new to Ethiopia and other notes. *Bulletin of the British Ornithologists' Club* 97: 4-9.
- Barnes, K. N. 2000. The Eskom Red Data Book of birds of South Africa, Lesotho and Swaziland. BirdLife South Africa, Johannesburg.
- Belik, V. P.; Lebedeva, E. A. 2004. International single-species action plan for the conservation of the Black-winged Pratincole *Glareola nordmanni*. AEWA Secretariat, Bonn, Germany.
- Belik, V.; Babich, M. V.; Korneev, P. I. 2000. [Catastrophic declines of Black-winged Pratincole (*Glareola nordmanni*) numbers north of the Caucasus]. *Information Materials of the Russian Wader Study Group* 13: 36-38.
- Berezovikov, N. N. 2002. [Decline in numbers of the Black-winged Pratincole in Eastern Kazakhstan]. *Information Materials of the Russian Wader Study Group* 15: 46-49.
- Bird, J. P., Martin, R., Akçakaya, H. R., Gilroy, J., Burfield, I. J., Garnett, S. G., Symes, A., Taylor, J., Sekercioglu, Ç. H. and Butchart, S. H. M. 2020. Generation lengths of the world's birds and their implications for extinction risk. *Conservation Biology* 34(5): 1252-1261. DOI: 10.1111/cobi.13486.
- BirdLife International. 2004. Birds in Europe: population estimates, trends and conservation status. BirdLife International, Cambridge, U.K.
- Cramp, S. and Simmons, K.E.L. 1983. Handbook of the birds of Europe, the Middle East and Africa. The birds of the western Palearctic vol. III: waders to gulls. Oxford University Press, Oxford.
- Dodman, T. 2002. Waterbird population estimates in Africa.
- Hayman, P., Marchant, J. and Prater, A.J. 1986. Shorebirds. Croom Helm, London.
- Hockey, P. A. R., Dean, W. R. J. & Ryan, P. G. 2005. Roberts Birds of Southern Africa. Trustees of the John Voelcker Bird Book Fund, Cape Town, South Africa.
- Hockey, P. and Douie, C. 1995. Struik Winchester, Cape Town, South Africa.
- Kamp, J.; Koshkin, M. A.; Sheldon, R. D. 2009. Population size, breeding performance and habitat use of the Black-winged Pratincole *Glareola nordmanni*. *Bird Conservation International* 19(2): 149-163.
- Kamp, J.; Koshkin, M. A.; Sheldon, R. D. 2009. Population size, breeding performance and habitat use of the Black-winged Pratincole *Glareola nordmanni*. *Bird Conservation International* 19(2): 149-163.
- Kamp, J.; Koshkin, M.; Sheldon, R. D. in prep. Population numbers, conservation status and habitat selection of the Black-winged Pratincole *Glareola nordmanni* in Kazakhstan.
- Karyakin, I. V.; Koslov, A. A. 1999. Stepnaya tirkushka (*Glareola nordmanni*).
- Maclean, G.L. 1996. Glareolidae (Courser and Pratincoles). In: J. del Hoyo, A. Elliott & J. Sargatal (eds), *Handbook of the birds of the world*, pp. 364-383. Lynx Edicions, Barcelona, Spain.
- Tucker, G.M. and Heath, M.F. 1994. Birds in Europe: Their Conservation Status. BirdLife International, Cambridge, U.K.
- University of Cape Town. 2006. Citizen science, again...: coordinated waterbird counts in South Africa. University of Cape Town, Cape Town, South Africa.
- del Hoyo, J., Elliott, A. and Sargatal, J. (eds). 1996. *Handbook of the Birds of the World, Vol. 3: Hoatzin to Auks*. Lynx Edicions, Barcelona, Spain.

du Plessis, G. J. 1995. Large aggregation of Black-winged Pratincoles *Glareola nordmanni* in the northern Orange Free State. *Ostrich* 66: 40-41.

du Plessis, G. J. 1995. Large aggregation of Black-winged Pratincoles *Glareola nordmanni* in the northern Orange Free State. *Ostrich* 66: 40-41.