

## *Oenanthe xanthopyrmyna* (Kurdish Wheatear)

### European Red List of Birds

### Supplementary Material

The European Union (EU28) Red List assessments were based principally on the official data reported by EU Member States to the European Commission under Article 12 of the Birds Directive in 2019-20. For the European Red List assessments, similar data were sourced from BirdLife Partners and other collaborating experts in other European countries and territories. For more information, see BirdLife International (2021).

#### Contents

- Reported national population sizes and trends
- Trend maps of reported national population data
- Sources of reported national population data
- Species factsheet bibliography

#### Recommended citation

BirdLife International (2021) European Red List of Birds. Luxembourg: Publications Office of the European Union.

#### Further information

<http://datazone.birdlife.org/info/euroredlist>  
<http://www.birdlife.org/europe-and-central-asia/european-red-list-birds-0>  
<http://www.iucnredlist.org/regions/europe>  
<http://ec.europa.eu/environment/nature/conservation/species/redlist/>

#### Data requests and feedback

To request access to these data in electronic format, provide new information, correct any errors or provide feedback, please email [science@birdlife.org](mailto:science@birdlife.org).

*Oenanthe xanthopyrma* (Kurdish Wheatear)

**Table 1.** Reported national breeding population size and trends in Europe<sup>1</sup>.

Country (or territory) <sup>2</sup>	Population estimate				Short-term population trend <sup>5</sup>				Long-term population trend <sup>5</sup>				Subspecific population (where relevant)
	Size (pairs) <sup>3</sup>	Europe (%)	Year(s)	Method <sup>4</sup>	Direction <sup>6</sup>	Magnitude (%) <sup>7</sup>	Year(s)	Method <sup>4</sup>	Direction <sup>6</sup>	Magnitude (%) <sup>7</sup>	Year(s)	Method <sup>4</sup>	
Azerbaijan	500–5000	69	1996-2019	expert	?		2013-2019	expert	?		1980-2019	expert	
Georgia	present	<1		deficient	?			deficient	?				
Turkey	500–1000	31	2002-2012	expert	?		2008-2019	deficient	?		1980-2013	deficient	
<b>Europe</b>	<b>1000–6000</b>	<b>100</b>											

<sup>1</sup> See 'Sources' at end of factsheet, and for more details on individual EU Member State reports, see the Article 12 reporting portal at <http://bd.eionet.europa.eu/article12/report>.

<sup>2</sup> The designation of geographical entities and the presentation of the material do not imply the expression of any opinion whatsoever on the part of IUCN or BirdLife International concerning the legal status of any country, territory or area, or of its authorities, or concerning the delimitation of its frontiers or boundaries.

<sup>3</sup> In the few cases where population size estimates were reported in units other than those specified, they were converted to the correct units using standard correction factors.

<sup>4</sup> The 'method used' (replacing the data 'quality' assessment in the 2015 European Red List) is reported as: a) Complete: complete survey or a statistically robust estimate; b) Partial: based mainly on extrapolation from a limited amount of data; c) Expert: based mainly on expert opinion with very limited data; d) Deficient: insufficient or no data available.

<sup>5</sup> The robustness of regional trends to the effects of any missing or incomplete data was tested using plausible scenarios, based on other sources of information, including any other reported information, recent national Red Lists, scientific literature, other publications and consultation with relevant experts.

<sup>6</sup> Trend directions are reported as: increasing (+); decreasing (-); stable (0); fluctuating (F); or unknown (?).

<sup>7</sup> Trend magnitudes are rounded to the nearest integer.

## Trend maps

A symbol appears in each country where the species occurs: the shape and colour of the symbol represent the population trend in that country, and the size of the symbol corresponds to the proportion of the European population occurring in that country.

### KEY


- |   |                                 |
|---|---------------------------------|
| ↑ Large increase (≥50%)                 | ↓ Large decrease (≥50%)         |
| ↑ Moderate increase (20–49%)            | ↓ Moderate decrease (20–49%)    |
| ↑ Small increase (<20%)                 | ↓ Small decrease (<20%)         |
| ↑ Increase of unknown magnitude         | ↓ Decrease of unknown magnitude |
| ■ Stable or fluctuating                 |                                 |
| □ Unknown                               |                                 |
| ○ Present (no population or trend data) |                                 |
| × Extinct since 1980                    |                                 |

Each symbol, with the exception of Present and Extinct, may occur in up to three different size classes, corresponding to the proportion of the European population occurring in that country.


- ↑ Large: ≥10% of the European population
- ↑ Medium: 1–9% of the European population
- ↑ Small: <1% of the European population

The designation of geographical entities and the presentation of the material do not imply the expression of any opinion whatsoever on the part of IUCN or BirdLife International concerning the legal status of any country, territory or area, or of its authorities, or concerning the delimitation of its frontiers or boundaries.

**Figure 1.** Breeding population sizes and short-term trends across Europe.



**Figure 2.** Breeding population sizes and long-term trends across Europe.



*Oenanthe xanthopyrna* (Kurdish Wheatear)

## Sources

### Azerbaijan

**Breeding population size:** BirdLife International 2004

**Breeding short-term trend:** AOS data base

**Breeding long-term trend:** AOS Data Base

### Georgia

**Breeding population size:** BirdLife International 2004

### Turkey

**Breeding population size:** Ferdi Akarsu, Güven Eken, Soner Bekir personal communication (2019), Kusbank Bird Database (Ebird) Kirwan G.M., Boyla K. A., Castell P., Demirci B., Özen M., Welch H., Marlow T., 2008, Birds of Turkey. Londra, Christopher Helm, 978-1-4081-0475-

## Bibliography

Bird, J. P., Martin, R., Akçakaya, H. R., Gilroy, J., Burfield, I. J., Garnett, S. G., Symes, A., Taylor, J., Sekercioglu, Ç. H. and Butchart, S. H. M. 2020. Generation lengths of the world's birds and their implications for extinction risk. *Conservation Biology* 34(5): 1252-1261. DOI: 10.1111/cobi.13486.

Collar, N.; de Juana, E. 2015. Chestnut-rumped Wheatear (*Oenanthe xanthopyrma*). In: del Hoyo, J.; Elliott, A.; Sargatal, J.; Christie, D. A.; de Juana, E. (ed.), *Handbook of the Birds of the World Alive*, Lynx Edicions, Barcelona, Spain.