



THE IUCN RED LIST
OF THREATENED SPECIES™



***Bombycilla garrulus* (Bohemian Waxwing)**

European Red List of Birds

Supplementary Material

The European Union (EU28) Red List assessments were based principally on the official data reported by EU Member States to the European Commission under Article 12 of the Birds Directive in 2019-20. For the European Red List assessments, similar data were sourced from BirdLife Partners and other collaborating experts in other European countries and territories. For more information, see BirdLife International (2021).

Contents

Reported national population sizes and trends
Trend maps of reported national population data
Sources of reported national population data
Species factsheet bibliography

Recommended citation

BirdLife International (2021) European Red List of Birds. Luxembourg: Publications Office of the European Union.

Further information

<http://datazone.birdlife.org/info/euroredlist>
<http://www.birdlife.org/europe-and-central-asia/european-red-list-birds-0>
<http://www.iucnredlist.org/regions/europe>
<http://ec.europa.eu/environment/nature/conservation/species/redlist/>

Data requests and feedback

To request access to these data in electronic format, provide new information, correct any errors or provide feedback, please email science@birdlife.org.

Bombycilla garrulus (Bohemian Waxwing)

Table 1. Reported national breeding population size and trends in Europe¹.

Country (or territory) ²	Population estimate				Short-term population trend ⁵				Long-term population trend ⁵				Subspecific population (where relevant)
	Size (pairs) ³	Europe (%)	Year(s)	Method ⁴	Direction ⁶	Magnitude (%) ⁷	Year(s)	Method ⁴	Direction ⁶	Magnitude (%) ⁷	Year(s)	Method ⁴	
Finland	52100–106000	8	2013-2018	complete	?	-2 to 70	2007-2018	complete	+	218 to 2110	1988-2018	complete	
Norway	100–2000	<1	2013-2018	expert	F		2013-2018	expert	F		1980-2018	partial	
Russia	500000–1000000	78	2006-2018	partial	F		2008-2019	expert	0		1980-2018	expert	
Sweden	98000–159000	14	2013-2018	partial	+	22 to 98	2007-2018	partial	+	50 to 150	1980-2018	partial	
EU28	150000–265000	22											
Europe	650000–1270000	100											

¹ See 'Sources' at end of factsheet, and for more details on individual EU Member State reports, see the Article 12 reporting portal at <http://bd.eionet.europa.eu/article12/report>.

² The designation of geographical entities and the presentation of the material do not imply the expression of any opinion whatsoever on the part of IUCN or BirdLife International concerning the legal status of any country, territory or area, or of its authorities, or concerning the delimitation of its frontiers or boundaries.

³ In the few cases where population size estimates were reported in units other than those specified, they were converted to the correct units using standard correction factors.

⁴ The 'method used' (replacing the data 'quality' assessment in the 2015 European Red List) is reported as: a) Complete: complete survey or a statistically robust estimate; b) Partial: based mainly on extrapolation from a limited amount of data; c) Expert: based mainly on expert opinion with very limited data; d) Defficient: insufficient or no data available.

⁵ The robustness of regional trends to the effects of any missing or incomplete data was tested using plausible scenarios, based on other sources of information, including any other reported information, recent national Red Lists, scientific literature, other publications and consultation with relevant experts.

⁶ Trend directions are reported as: increasing (+); decreasing (-); stable (0); fluctuating (F); or unknown (?).

⁷ Trend magnitudes are rounded to the nearest integer.

Bombycilla garrulus (Bohemian Waxwing)

Table 2. Reported national wintering population sizes and trends in Europe¹. Note that some countries within the species' wintering range did not report any data, and that only minimum totals are presented, to avoid double-counting of birds moving between countries.

Country (or territory) ²	Population estimate				Short-term population trend ⁵				Long-term population trend ⁵				Subspecific population (where relevant)
	Size (individuals) ³	Europe (%)	Year(s)	Method ⁴	Direction ⁶	Magnitude (%) ⁷	Year(s)	Method ⁴	Direction ⁶	Magnitude (%) ⁷	Year(s)	Method ⁴	
Czechia	100–50000	69	2014-2018	expert	F		2007-2018	expert	F		1982-2018	expert	
Serbia	1000	31	2013-2018	partial	F		2013-2018	complete	F		1980-2018	complete	
EU28	100–50000	69											
Europe	100–51000	100											

¹ See 'Sources' at end of factsheet, and for more details on individual EU Member State reports, see the Article 12 reporting portal at <http://bd.eionet.europa.eu/article12/report>.

² The designation of geographical entities and the presentation of the material do not imply the expression of any opinion whatsoever on the part of IUCN or BirdLife International concerning the legal status of any country, territory or area, or of its authorities, or concerning the delimitation of its frontiers or boundaries.

³ In the few cases where population size estimates were reported in units other than those specified, they were converted to the correct units using standard correction factors.

⁴ The 'method used' (replacing the data 'quality' assessment in the 2015 European Red List) is reported as: a) Complete: complete survey or a statistically robust estimate; b) Partial: based mainly on extrapolation from a limited amount of data; c) Expert: based mainly on expert opinion with very limited data; d) Defficient: insufficient or no data available.

⁵ The robustness of regional trends to the effects of any missing or incomplete data was tested using plausible scenarios, based on other sources of information, including any other reported information, recent national Red Lists, scientific literature, other publications and consultation with relevant experts.

⁶ Trend directions are reported as: increasing (+); decreasing (-); stable (0); fluctuating (F); or unknown (?).

⁷ Trend magnitudes are rounded to the nearest integer.

Trend maps

A symbol appears in each country where the species occurs: the shape and colour of the symbol represent the population trend in that country, and the size of the symbol corresponds to the proportion of the European population occurring in that country.

KEY

- | | |
|---|---------------------------------|
| ↑ Large increase (≥50%) | ↓ Large decrease (≥50%) |
| ↑ Moderate increase (20–49%) | ↓ Moderate decrease (20–49%) |
| ↑ Small increase (<20%) | ↓ Small decrease (<20%) |
| ↑ Increase of unknown magnitude | ↓ Decrease of unknown magnitude |
| ■ Stable or fluctuating | |
| □ Unknown | |
| ○ Present (no population or trend data) | |
| × Extinct since 1980 | |

Each symbol, with the exception of Present and Extinct, may occur in up to three different size classes, corresponding to the proportion of the European population occurring in that country.

- ↑ Large: ≥10% of the European population
- ↑ Medium: 1–9% of the European population
- ↑ Small: <1% of the European population

The designation of geographical entities and the presentation of the material do not imply the expression of any opinion whatsoever on the part of IUCN or BirdLife International concerning the legal status of any country, territory or area, or of its authorities, or concerning the delimitation of its frontiers or boundaries.

Figure 1. Breeding population sizes and short-term trends across Europe.




Figure 2. Breeding population sizes and long-term trends across Europe.




Figure 3. Reported wintering population sizes and short-term trends across Europe. Note that some countries within the species' wintering range did not report any data.





Figure 4. Reported wintering population sizes and long-term trends across Europe. Note that some countries within the species' wintering range did not report any data.



Sources

Czechia

Winter population size: expert opinion
Winter short-term trend: expert opinion
Winter long-term trend: expert opinion

Finland

Breeding population size: Lehtinen, A., Below, A., Jukarainen, A., Laaksonen, T., Lehtinen, T., Mikkola-Roos, M., Pessa, J., Rajasärkkä, A., Rusanen, P., Sirkiä, P., Tiainen, J. & Valkama, J. 2019: Suomen lintujen pesimäkantojen koot. – Linnut-vuosikirja 2018: 38-45.
Breeding short-term trend: Väisänen, R. A., Lehtinen, A. & Sirkiä, P. 2018: Suomen pesivän maalinuston kannanvaihtelut 1975-2017. Linnut-vuosikirja 2017: 16–31.
Breeding long-term trend: Väisänen, R. A., Lehtinen, A. & Sirkiä, P. 2018: Suomen pesivän maalinuston kannanvaihtelut 1975-2017. Linnut-vuosikirja 2017: 16–31.

Norway

Breeding population size: Shimmings P. & Øien, I.J. 2015. Bestandsestimater og trender for norske hekkefugler. NOF-rapport 2015-2.
Breeding short-term trend: Shimmings P. & Øien, I.J. 2015. Bestandsestimater og trender for norske hekkefugler. NOF-rapport 2015-2.
Breeding long-term trend: Shimmings, P. & Øien, I.J. 2015. Bestandsestimater for norske hekkefugler. NOF Rapport 2-2015. 268 pp.

Russia

Breeding population size: Voltz & Kalyakin 2013-2019; Database of the project on Atlas of breeding birds of European Russia
Breeding short-term trend: Yakovleva unpublished. kivach-bird@rambler.ru Preobrazhenskaya unpublished. voop21@rambler.ru
Breeding long-term trend: Yakovleva unpublished. kivach-bird@rambler.ru Preobrazhenskaya unpublished. voop21@rambler.ru

Serbia

Winter population size: Bioras database http://www.bioras.petnica.rs/home.php
Winter short-term trend: Bioras database http://www.bioras.petnica.rs/home.php
Winter long-term trend: Bioras database http://www.bioras.petnica.rs/home.php

Sweden

Breeding population size: Ottosson, U., Ottvall, R., Elmqvist, J., Green, M., Gustafsson, R., Haas, F., Holmqvist, N., Lindström, Å., Nilsson, L., Svensson, M., Svensson, S. & Tjernberg, M. 2012. Fåglarna i Sverige – antal och förekomst. SOF, Halmstad. Swedish Bird Survey. BirdLife Sverige, Annual Bird reports.
Breeding short-term trend: Svensk fågeltaxering - Swedish Bird Survey
Breeding long-term trend: Svensk fågeltaxering - Swedish Bird Survey

Bibliography

- Bird, J. P., Martin, R., Akçakaya, H. R., Gilroy, J., Burfield, I. J., Garnett, S. G., Symes, A., Taylor, J., Sekercioglu, Ç. H. and Butchart, S. H. M. 2020. Generation lengths of the world's birds and their implications for extinction risk. *Conservation Biology* 34(5): 1252-1261. DOI: 10.1111/cobi.13486.
- Brazil, M. 2009. *Birds of East Asia: eastern China, Taiwan, Korea, Japan, eastern Russia*. Christopher Helm, London.
- Mountjoy, J. 2018. Bohemian Waxwing (*Bombycilla garrulus*). In: del Hoyo, J.; Elliott, A.; Sargatal, J.; Christie, D.A.; de Juana, E. (ed.), *Handbook of the Birds of the World Alive*, Lynx Edicions, Barcelona.
- Rich, T.D., Beardmore, C.J., Berlanga, H., Blancher, P.J., Bradstreet, M.S.W., Butcher, G.S., Demarest, D.W., Dunn, E.H., Hunter, W.C., Inigo-Elias, E.E., Martell, A.M., Panjabi, A.O., Pashley, D.N., Rosenberg, K.V., Rustay, C.M., Wendt, J.S. and Will, T.C. 2004. *Partners in flight: North American landbird conservation plan*. Cornell Lab of Ornithology, Ithaca, NY.
- Töpfer, T. 2010. Suspected road salt poisoning in Bohemian waxwings *Bombycilla garrulus* (Aves: Passeriformes: Bombycillidae). *Vertebrate Zoology* 60(2): 171-174.