



THE IUCN RED LIST
OF THREATENED SPECIES™



***Cinclus cinclus* (White-throated Dipper)**

European Red List of Birds

Supplementary Material

The European Union (EU28) Red List assessments were based principally on the official data reported by EU Member States to the European Commission under Article 12 of the Birds Directive in 2019-20. For the European Red List assessments, similar data were sourced from BirdLife Partners and other collaborating experts in other European countries and territories. For more information, see BirdLife International (2021).

Contents

Reported national population sizes and trends
Trend maps of reported national population data
Sources of reported national population data
Species factsheet bibliography

Recommended citation

BirdLife International (2021) European Red List of Birds. Luxembourg: Publications Office of the European Union.

Further information

<http://datazone.birdlife.org/info/euroredlist>
<http://www.birdlife.org/europe-and-central-asia/european-red-list-birds-0>
<http://www.iucnredlist.org/regions/europe>
<http://ec.europa.eu/environment/nature/conservation/species/redlist/>

Data requests and feedback

To request access to these data in electronic format, provide new information, correct any errors or provide feedback, please email science@birdlife.org.

Cinclus cinclus (White-throated Dipper)

Table 1. Reported national breeding population size and trends in Europe¹.

Country (or territory) ²	Population estimate				Short-term population trend ⁵				Long-term population trend ⁵				Subspecific population (where relevant)
	Size (pairs) ³	Europe (%)	Year(s)	Method ⁴	Direction ⁶	Magnitude (%) ⁷	Year(s)	Method ⁴	Direction ⁶	Magnitude (%) ⁷	Year(s)	Method ⁴	
Albania	300–630	<1	2007-2018	partial	?	-21 to 50	2007-2018	partial	-	-37 to 50	1980-2018	expert	
Andorra	100–240	<1	2014-2017	partial	-		2011-2018	complete	?				
Armenia	1400–2000	1	2013-2018	complete	0		2007-2018		0		2003-2018	partial	
Austria	6000–9000	4	2013-2018	partial	0		2007-2018	partial	?		1981-2018	deficient	
Azerbaijan	1000–10000	2	1996-2019	expert	?		2013-2019	expert	?		1980-2019	expert	
Belgium	740–1100	<1	2013-2018	partial	0	-18 to 17	2008-2018	partial	0	0 to 42	1973-2018	partial	
Bosnia & HG	1000–2000	<1	2015-2018	complete	?	-10 to 10	2007-2018	complete	?		1980-2018	deficient	
Bulgaria	8000–11000	6	2005-2018	partial	0	0 to 5	2000-2018	expert	0	0 to 5	1980-2018	expert	
Croatia	1000	<1	2014-2014	expert	?		2007-2018	deficient	?		1980-2018	deficient	
Czechia	1000–2000	<1	2014-2017	complete	?		2007-2018	deficient	?		1980-2018	deficient	
Denmark	4	<1	2017	complete	-		2006-2017	expert	?		1980-2017	partial	
Estonia	3–10	<1	2013-2017	expert	?	20 to 50	2006-2017	expert	?		1980-2017	deficient	
Finland	200–300	<1	2013-2018	partial	?		2007-2018	deficient	?		1980-2018	deficient	
France	15000–24000	11	2013-2018	partial	?		2007-2017	deficient	0	0	1990-2017	expert	
Georgia	1600–16600	3	2013-2017	partial	?			deficient	?				
Germany	11000–19500	9	2011-2016	expert	0		2004-2016	expert	+		1980-2016	expert	
Greece	500–2000	<1	2015	partial	0		2007-2018	partial	0		1980-2018	partial	
Hungary	0–4	<1	2015-2017	complete	F		2007-2017	partial	-		1980-2017	complete	
Rep. Ireland	2000–6100	2	2008-2011	partial	+		1991-2011	expert	-		1972-2011	expert	
Italy	4000–8000	3	2013-2018	expert	?		2007-2018	deficient	0		1993-2018	expert	
Kosovo	600–800	<1	2007-2019	partial	-		2007-2018	partial	0		1990-2018	partial	
Latvia	1–10	<1	2013-2018	complete	+	36 to 166	2012-2018	complete	+	0 to 110	1991-2017	partial	
Luxembourg	100–200	<1	2013-2018	partial	-	-20 to -10	2007-2018	partial	-	-30 to -10	1980-2018	expert	
North Macedonia	1000–3000	1	2014-2019	expert	0		2007-2018	expert	?		1980-2019		
Montenegro	500–1500	<1	2002-2012	expert	0		2007-2018	expert	?				
Norway	10000–40000	12	2013-2018	expert	?		2013-2018	deficient	?		1980-2018	partial	
Poland	1900–3000	1	2013-2014	partial	?		2007-2018	deficient	?		1980-2018	deficient	
Portugal	500–2500	<1	2013-2018	partial	-		2007-2018	partial	?		1980-2018	deficient	
Romania	2600–26000	5	2013-2018	expert	-	-20 to -1	2007-2018	expert	-	-50 to -10	1980-2018	expert	
Russia	7900–14000	6	2008-2018	partial	-	-49 to -30	2008-2019	expert	-	-79 to -50	1980-2018	expert	
Serbia	2400–3400	2	2013-2018	partial	-	-29 to -10	2007-2018	complete	0	0	1980-2018	complete	
Slovakia	2000–3000	2	2013-2018	partial	0		2007-2018	partial	0		1980-2018	partial	

Cinclus cinclus (White-throated Dipper)

Table 1. Reported national breeding population size and trends in Europe¹.

Country (or territory) ²	Population estimate				Short-term population trend ⁵				Long-term population trend ⁵				Subspecific population (where relevant)
	Size (pairs) ³	Europe (%)	Year(s)	Method ⁴	Direction ⁶	Magnitude (%) ⁷	Year(s)	Method ⁴	Direction ⁶	Magnitude (%) ⁷	Year(s)	Method ⁴	
Slovenia	1000–3000	1	2002-2017	complete	?		2008-2018	deficient	?		1980-2018	deficient	
Spain	3300–3400	2	1998-2018	complete	?	-3 to 9	2007-2018	complete	?		1980-2018	deficient	
Sweden	5700–15000	6	2013-2018	partial	?	-48 to 417	2007-2018	partial	0	-35 to 8	1980-2018	partial	
Switzerland	6000–8000	4	2013–2016	partial	0	-39 to 39	2007-2018	complete	+	3 to 78	1990-2018	complete	
Turkey	2800–8400	3	2002-2012	deficient	?		2008-2019	deficient	?		1980-2013	deficient	
Ukraine	1200–1800	<1	2015-2017	partial	F	5 to 10	2007-2019	expert	F	10 to 15	1980-2019	expert	
United Kingdom	6900–20800	7	2016	partial	0		2004-2016	complete	0		1980-2016	complete	
EU28	73400–161000	63											
Europe	111000–274000	100											

¹ See 'Sources' at end of factsheet, and for more details on individual EU Member State reports, see the Article 12 reporting portal at <http://bd.eionet.europa.eu/article12/report>.

² The designation of geographical entities and the presentation of the material do not imply the expression of any opinion whatsoever on the part of IUCN or BirdLife International concerning the legal status of any country, territory or area, or of its authorities, or concerning the delimitation of its frontiers or boundaries.

³ In the few cases where population size estimates were reported in units other than those specified, they were converted to the correct units using standard correction factors.

⁴ The 'method used' (replacing the data 'quality' assessment in the 2015 European Red List) is reported as: a) Complete: complete survey or a statistically robust estimate; b) Partial: based mainly on extrapolation from a limited amount of data; c) Expert: based mainly on expert opinion with very limited data; d) Defficient: insufficient or no data available.

⁵ The robustness of regional trends to the effects of any missing or incomplete data was tested using plausible scenarios, based on other sources of information, including any other reported information, recent national Red Lists, scientific literature, other publications and consultation with relevant experts.

⁶ Trend directions are reported as: increasing (+); decreasing (-); stable (0); fluctuating (F); or unknown (?).

⁷ Trend magnitudes are rounded to the nearest integer.

Cinclus cinclus (White-throated Dipper)

Table 2. Reported national wintering population sizes and trends in Europe¹. Note that some countries within the species' wintering range did not report any data, and that only minimum totals are presented, to avoid double-counting of birds moving between countries.

Country (or territory) ²	Population estimate				Short-term population trend ⁵				Long-term population trend ⁵				Subspecific population (where relevant)
	Size (individuals) ³	Europe (%)	Year(s)	Method ⁴	Direction ⁶	Magnitude (%) ⁷	Year(s)	Method ⁴	Direction ⁶	Magnitude (%) ⁷	Year(s)	Method ⁴	
Bosnia & HG	3000–4000	38	2015-2018	complete	?		2007-2018	deficient	?		1980-2018	deficient	
Czechia	2500–5000	38	2015-2019	complete	+		2008-2019	complete	+		1980-2019	complete	
Serbia	1000–5000	24	2013-2018	partial	?	-10 to 10	2013-2018	partial	?	-10 to 10	1980-2018	expert	
EU28	2500–5000	38											
Europe	6500–14000	100											

¹ See 'Sources' at end of factsheet, and for more details on individual EU Member State reports, see the Article 12 reporting portal at <http://bd.eionet.europa.eu/article12/report>.

² The designation of geographical entities and the presentation of the material do not imply the expression of any opinion whatsoever on the part of IUCN or BirdLife International concerning the legal status of any country, territory or area, or of its authorities, or concerning the delimitation of its frontiers or boundaries.

³ In the few cases where population size estimates were reported in units other than those specified, they were converted to the correct units using standard correction factors.

⁴ The 'method used' (replacing the data 'quality' assessment in the 2015 European Red List) is reported as: a) Complete: complete survey or a statistically robust estimate; b) Partial: based mainly on extrapolation from a limited amount of data; c) Expert: based mainly on expert opinion with very limited data; d) Defficient: insufficient or no data available.

⁵ The robustness of regional trends to the effects of any missing or incomplete data was tested using plausible scenarios, based on other sources of information, including any other reported information, recent national Red Lists, scientific literature, other publications and consultation with relevant experts.

⁶ Trend directions are reported as: increasing (+); decreasing (-); stable (0); fluctuating (F); or unknown (?).

⁷ Trend magnitudes are rounded to the nearest integer.

Trend maps

A symbol appears in each country where the species occurs: the shape and colour of the symbol represent the population trend in that country, and the size of the symbol corresponds to the proportion of the European population occurring in that country.

KEY

- | | |
|---|---------------------------------|
| ↑ Large increase (≥50%) | ↓ Large decrease (≥50%) |
| ↑ Moderate increase (20–49%) | ↓ Moderate decrease (20–49%) |
| ↑ Small increase (<20%) | ↓ Small decrease (<20%) |
| ↑ Increase of unknown magnitude | ↓ Decrease of unknown magnitude |
| ■ Stable or fluctuating | |
| □ Unknown | |
| ○ Present (no population or trend data) | |
| × Extinct since 1980 | |

Each symbol, with the exception of Present and Extinct, may occur in up to three different size classes, corresponding to the proportion of the European population occurring in that country.

- ↑ Large: ≥10% of the European population
- ↑ Medium: 1–9% of the European population
- ↑ Small: <1% of the European population

The designation of geographical entities and the presentation of the material do not imply the expression of any opinion whatsoever on the part of IUCN or BirdLife International concerning the legal status of any country, territory or area, or of its authorities, or concerning the delimitation of its frontiers or boundaries.

Figure 1. Breeding population sizes and short-term trends across Europe.




Figure 2. Breeding population sizes and long-term trends across Europe.




Figure 3. Reported wintering population sizes and short-term trends across Europe. Note that some countries within the species' wintering range did not report any data.




Figure 4. Reported wintering population sizes and long-term trends across Europe. Note that some countries within the species' wintering range did not report any data.



Cinclus cinclus (White-throated Dipper)

Sources

Albania

Breeding population size: Bino & Xeka pers. obs.
Breeding short-term trend: Bino & Xeka pers. obs.
Breeding long-term trend: Bino pers. obs.

Andorra

Breeding population size: Fieldwork EBBA2, published at "Guia dels ocells d'Andorra. J. Nicolau & C. Pladevall, 2018"
Breeding short-term trend: Common Bird Monitoring Scheme of Andorra (SOCA)

Armenia

Breeding population size: TSE NGO National Bird Monitoring data.
Breeding short-term trend: TSE (2020) The Atlas of the Breeding Birds in Armenia. In preparation.
Breeding long-term trend: TSE (2020) The Atlas of the Breeding Birds in Armenia. In preparation.

Austria

Breeding population size: BirdLife Austria, estimate based on a sample of breeding densities from different sites and habitats
Breeding short-term trend: BirdLife Austria, unpublished data from www.ornitho.at ; BirdLife Austria, unpublished archive data
Breeding long-term trend: BirdLife Austria, unpublished archive data

Azerbaijan

Breeding population size: BirdLife International 2004
Breeding short-term trend: AOS data base
Breeding long-term trend: AOS Data Base

Belgium

Breeding population size: Vermeersch G. et al. (2018, in press). Broedvogels in Vlaanderen in de periode 2013-2018. Rapporten van het Instituut voor Natuur- en Bosonderzoek (INBO), Brussel. / Paquet, J-Y., Anselin, A., Vermeersch, G., Derouaux, A., Devos, K. (2019, in prep.). Contribution of Belgium to EBCC European Breeding Bird Atlas 2. Internal Report.
Breeding short-term trend: Vermeersch G. et al. (2018, in press). Broedvogels in Vlaanderen in de periode 2013-2018. Rapporten van het Instituut voor Natuur- en Bosonderzoek (INBO), Brussel. / Paquet, J-Y., Anselin, A., Vermeersch, G., Derouaux, A., Devos, K. (2019, in prep.). Contribution of Belgium to EBCC European Breeding Bird Atlas 2. Internal Report.
Breeding long-term trend: Vermeersch G. et al. (2018, in press). Broedvogels in Vlaanderen in de periode 2013-2018. Rapporten van het Instituut voor Natuur- en Bosonderzoek (INBO), Brussel. / Paquet, J-Y., Anselin, A., Vermeersch, G., Derouaux, A., Devos, K. (2019, in prep.). Contribution of Belgium to EBCC European Breeding Bird Atlas 2. Internal Report.

Bosnia and Herzegovina

Breeding population size: Based on data for EBBA2
Breeding short-term trend: more individual articles e.g published in magazine Bilten mreže posmatrača ptica u Bosni i Hercegovini-see https://ptice.ba/bs/category/bilteni_/ , individual reports (e.g. for EBBA2, projects etc)
Winter population size: based on IWC reports-all reports published in magazine Bilten mreže posmatrača ptica (www.ptice.ba)
Winter short-term trend: based on IWC reports-all reports published in magazine Bilten mreže posmatrača ptica (www.ptice.ba)
Winter long-term trend: There are no qualitative data before 2005 to make estimates

Bulgaria

Breeding population size: Iankov, P. (ed.) 2007 Atlas of Breeding Birds in Bulgaria. Bulgarian Society for the Protection of Birds, Conservation Series, Book 10, Sofia, BSPB, 679 p.; National Art. 12 reporting database 2013-2018; Nankinov, D. 2009. Studies on Fauna of Bulgaria, Birds - Aves, Passeriformes, Sofia, ETO, 407 p. (in Bulgarian) BSPB Bird Database
Breeding short-term trend: Iankov, P. (ed.) 2007 Atlas of Breeding Birds in Bulgaria. Bulgarian Society for the Protection of Birds, Conservation Series, Book 10, Sofia, BSPB, 679 p.; National Art. 12 reporting database 2013-2018; Nankinov, D. 2009. Studies on Fauna of Bulgaria, Birds - Aves, Passeriformes, Sofia, ETO, 407 p. (in Bulgarian) BSPB Bird Database
Breeding long-term trend: Iankov, P. (ed.) 2007 Atlas of Breeding Birds in Bulgaria. Bulgarian Society for the Protection of Birds, Conservation Series, Book 10, Sofia, BSPB, 679 p. Nankinov, D. 2009. Studies on Fauna of Bulgaria, Birds - Aves, Passeriformes, Sofia, ETO, 407 p. (in Bulgarian) BSPB Bird Database

Croatia

Breeding population size: Dumbović Mazal V., Pintar V., Zadavec M. (2019): Prvo izvješće o brojnosti i rasprostranjenosti ptica u Hrvatskoj sukladno odredbama Direktive o pticama.
Breeding short-term trend: no data available
Breeding long-term trend: no data available

Czechia

Breeding population size: Šťastný et Bejček in prep. - Atlas hnízdního rozšíření ptáků ČR 2014-2017
--

Cinclus cinclus (White-throated Dipper)

Czechia

Breeding short-term trend: expert opinion
Breeding long-term trend: expert opinion
Winter population size: Waterbird numbers were recorded in mid-January by regular citizen-science monitoring programme - the International Waterbird Census (IWC) – see Gilissen et al. 2002, Wetlands International 2006, Wetlands International 2019. Hundreds of volunteer birdwatchers conduct the mid-January counts on predetermined dates and sites each year, aiming to maximize synchrony (Gilissen et al. 2002, Musil et al. 2011, Musilová et al. 2014). The estimation of population size was calculated by Trends and Indices for Monitoring data (TRIM) software (Statistics Netherlands version 3.52, Pannekoek and Van Strien, 2005). 'Time Totals' values of the data (i.e. the actual count values plus the numbers of birds estimated by the TRIM software) for all 1155 sites included in the analysis were used to generate total estimates of the range of numbers of the waterbird species wintering in the Czech Republic between 2015 and 2019. We use the range (min–max) of population estimates due to the effect of between-year variation in numbers because of variable climatic conditions (Musil et al. 2008, Musilová et al. 2018). Gilissen N, Haanstra L, Delany S, Boere G, Hagemeyer W (2002) Numbers and distribution of wintering waterbirds in the Western Palearctic and Southwest Asia in 1987, 1988 and 1999. Results from the International Waterbird Census. Wetlands International Global Series No. 11, Wetlands International, Wageningen, The Netherlands. Musil P, Darolová A, Jureček J, Musilová Z, Podhrázký M, Slabeyová K (2008) The long-term trends in numbers of wintering geese in the Czech Republic and Slovakia in 1991–2007. Tichodroma 20: 61–67. Musil P, Musilová Z, Fuchs R, Poláková S (2011) Long-term changes in numbers and distribution of wintering waterbirds in the Czech Republic, 1966–2008. Bird Study 58: 450–460. Musilová Z, Musil P, Zouhar J, Adam M (2018) Changes in habitat suitability influence non-breeding distribution of waterbirds in central Europe. Ibis: 160: 582–596. Musilová Z, Musil P, Zouhar J, Bejček V, Šťastný K, Hudec K (2014) Numbers of wintering waterbirds in the Czech Republic: long-term and spatial-scale approaches to assess population size. Bird Study 61: 321–331.
Winter short-term trend: Waterbird numbers were recorded in mid-January by regular citizen-science monitoring programme - the International Waterbird Census (IWC) – see Gilissen et al. 2002, Wetlands International 2006, Wetlands International 2019. Hundreds of volunteer birdwatchers conduct the mid-January counts on predetermined dates and sites each year, aiming to maximize synchrony (Gilissen et al. 2002, Musil et al. 2011, Musilová et al. 2014). The individual species trends in numbers was calculated by Trends and Indices for Monitoring data (TRIM) software (Statistics Netherlands version 3.52, Pannekoek and Van Strien, 2005). The additive slope (i.e. the change in indices from one year to the next) was used to estimate the Czech trend, see also Fouque et al. (2009), Musil et al. (2011), Musilová et al. (2015), Musilová et al. (2018 a, b). Fouque C, Guillemain M, Schricke V (2009) Trends in the numbers of Coot Fulica atra and wildfowl Anatidae wintering in France and their relationship with hunting activity at wetland sites. Wildfowl. Special Issue 2: 42–59. Gilissen N, Haanstra L, Delany S, Boere G, Hagemeyer W (2002) Numbers and distribution of wintering waterbirds in the Western Palearctic and Southwest Asia in 1987, 1988 and 1999. Results from the International Waterbird Census. Wetlands International Global Series No. 11, Wetlands International, Wageningen, The Netherlands. Musil P, Musilová Z, Fuchs R, Poláková S (2011) Long-term changes in numbers and distribution of wintering waterbirds in the Czech Republic, 1966–2008. Bird Study 58: 450–460. Musilová Z, Musil P, Zouhar J, Adam M (2018a) Changes in habitat suitability influence non-breeding distribution of waterbirds in central Europe. Ibis: 160: 582–596. Musilová Z, Musil P, Zouhar J, Adam M, Bejček V (2018b) Importance of Natura 2000 sites for wintering waterbirds: Low preference, species' distribution changes and carrying capacity of Natura 2000 could fail to protect the species. Biological Conservation 228: 79–88. Musilová Z, Musil P, Zouhar J, Bejček V, Šťastný K, Hudec K (2014) Numbers of wintering waterbirds in the Czech Republic: long-term and spatial-scale approaches to assess population size. Bird Study 61: 321–331. Musilová Z, Musil P, Zouhar J, Romportl D (2015) Long-term trends, total numbers and species richness of increasing waterbird populations at sites on the edge of their winter range: cold-weather refuge sites are more important than protected sites. J Ornithol: 1–10. Pannekoek J, Van Strien AJ (2005) TRIM 3 Manual (TRends and Indices for Monitoring Data). Statistics Netherlands, Voorburg, The Netherlands. Wetlands International (2006) Waterbird population estimates. Fourth Edition. Wetlands International, Wageningen, The Netherlands. Wetlands International (2019) Waterbird Population Estimates. Available at: wpe.wetlands.org (accessed 10 March 2019).
Winter long-term trend: Waterbird numbers were recorded in mid-January by regular citizen-science monitoring programme - the International Waterbird Census (IWC) – see Gilissen et al. 2002, Wetlands International 2006, Wetlands International 2019. Hundreds of volunteer birdwatchers conduct the mid-January counts on predetermined dates and sites each year, aiming to maximize synchrony (Gilissen et al. 2002, Musil et al. 2011, Musilová et al. 2014). The individual species trends in numbers was calculated by Trends and Indices for Monitoring data (TRIM) software (Statistics Netherlands version 3.52, Pannekoek and Van Strien, 2005). The additive slope (i.e. the change in indices from one year to the next) was used to estimate the Czech trend, see also Fouque et al. (2009), Musil et al. (2011), Musilová et al. (2015), Musilová et al. (2018 a, b). Fouque C, Guillemain M, Schricke V (2009) Trends in the numbers of Coot Fulica atra and wildfowl Anatidae wintering in France and their relationship with hunting activity at wetland sites. Wildfowl. Special Issue 2: 42–59. Gilissen N, Haanstra L, Delany S, Boere G, Hagemeyer W (2002) Numbers and distribution of wintering waterbirds in the Western Palearctic and Southwest Asia in 1987, 1988 and 1999. Results from the International Waterbird Census. Wetlands International Global Series No. 11, Wetlands International, Wageningen, The Netherlands. Musil P, Musilová Z, Fuchs R, Poláková S (2011) Long-term changes in numbers and distribution of wintering waterbirds in the Czech Republic, 1966–2008. Bird Study 58: 450–460. Musilová Z, Musil P, Zouhar J, Adam M (2018a) Changes in habitat suitability influence non-breeding distribution of waterbirds in central Europe. Ibis: 160: 582–596. Musilová Z, Musil P, Zouhar J, Adam M, Bejček V (2018b) Importance of Natura 2000 sites for wintering waterbirds: Low preference, species' distribution changes and carrying capacity of Natura 2000 could fail to protect the species. Biological Conservation 228: 79–88. Musilová Z, Musil P, Zouhar J, Bejček V, Šťastný K, Hudec K (2014) Numbers of wintering waterbirds in the Czech Republic: long-term and spatial-scale approaches to assess population size. Bird Study 61: 321–331. Musilová Z, Musil P, Zouhar J, Romportl D (2015) Long-term trends, total numbers and species richness of increasing waterbird populations at sites on the edge of their winter range: cold-weather refuge sites are more important than protected sites. J Ornithol: 1–10. Pannekoek J, Van Strien AJ (2005) TRIM 3 Manual (TRends and Indices for Monitoring Data). Statistics Netherlands, Voorburg, The Netherlands. Wetlands International (2006) Waterbird population estimates. Fourth Edition. Wetlands International, Wageningen, The Netherlands. Wetlands International (2019) Waterbird Population Estimates. Available at: wpe.wetlands.org (accessed 10 March 2019).

Denmark

Breeding population size: www.dofbasen.dk & Nyegaard, T. et al., Truede og sjældne ynglefugle i Danmark 1998-2012, Dansk Ornitologisk Forenings Tidsskrift 108, nr 1, 2014 & Atlas III 2014-2017 (www.dofbasen.dk/atlas) & DOF BirdLifeDK Fugleåret 2006-2017 &
Breeding short-term trend: www.dofbasen.dk & Nyegaard, T. et al., Truede og sjældne ynglefugle i Danmark 1998-2012, Dansk Ornitologisk Forenings Tidsskrift 108, nr 1, 2014 & Atlas III 2014-2017 (www.dofbasen.dk/atlas) & DOF BirdLifeDK Fugleåret 2006-2017
Breeding long-term trend: www.dofbasen.dk & Nyegaard, T. et al., Truede og sjældne ynglefugle i Danmark 1998-2012, Dansk Ornitologisk Forenings Tidsskrift 108, nr 1, 2014 & Atlas III 2014-2017 (www.dofbasen.dk/atlas) & DOF BirdLifeDK Fugleåret 2006-2017

Estonia

Breeding population size: Estonian Working Group on Bird Status and Numbers
Breeding short-term trend: Estonian Working Group on Bird Status and Numbers
Breeding long-term trend: Estonian Working Group on Bird Status and Numbers

Finland

Breeding population size: Lehikoinen, A., Below, A., Jukarainen, A., Laaksonen, T., Lehtiniemi, T., Mikkola-Roos, M., Pessa, J., Rajasärkkä, A., Rusanen, P., Sirkiä, P., Tiainen, J. & Valkama, J. 2019: Suomen lintujen pesimäkantojen koot. – Linnut-vuosikirja 2018: 38-45.
--

Cinclus cinclus (White-throated Dipper)

France

Breeding population size: Issa N. & Muller Y. 2015. Atlas des oiseaux nicheurs de France métropolitaine. , LPO/SEOF/MNHN/Delachaux et Niestlé, Paris

Breeding long-term trend: Roché J. & Faivre B. 2013. Suivi quantitatif de l'avifaune nicheuse de la Loire et de l'Allier de 1990 à 2012. Université de Bourgogne, 112 p.

Georgia

Breeding population size: EBBA Georgia, prepared by Sabuko-Society for nature conservation, Ilia state university, NGO "psovi".

Germany

Breeding population size: Gerlach et al. (in Vorb.): Vögel in Deutschland – 2019. Dachverband Deutscher Avifaunisten, Bundesamt für Naturschutz und Länderarbeitsgemeinschaft der Vogelschutzwarten, Münster.

Breeding short-term trend: Gerlach et al. (in Vorb.): Vögel in Deutschland – 2019. Dachverband Deutscher Avifaunisten, Bundesamt für Naturschutz und Länderarbeitsgemeinschaft der Vogelschutzwarten, Münster.

Breeding long-term trend: Gerlach et al. (in Vorb.): Vögel in Deutschland – 2019. Dachverband Deutscher Avifaunisten, Bundesamt für Naturschutz und Länderarbeitsgemeinschaft der Vogelschutzwarten, Münster.

Greece

Breeding population size: 1) BirdLife International (2004) Birds in Europe: Population estimates, trends and conservation status. Cambridge, UK: Birdlife International (Birdlife Conservation Series No. 12).

Breeding short-term trend: 1) BirdLife International (2004) Birds in Europe: Population estimates, trends and conservation status. Cambridge, UK: Birdlife International (Birdlife Conservation Series No. 12).

Breeding long-term trend: 1) Handrinos, G., & Akriotis, T., (1997) The birds of Greece. C. Helm, A & C Black, London. 2) BirdLife International (2004) Birds in Europe: Population estimates, trends and conservation status. Cambridge, UK: Birdlife International (Birdlife Conservation Series No. 12).

Hungary

Breeding population size: National park directorates' databases (Annual survey of colonially breeding and strictly protected bird species) <http://map.mme.hu/maps/map2/www.birding.hu>

Breeding short-term trend: National park directorates' databases (Annual survey of colonially breeding and strictly protected bird species) <http://map.mme.hu/maps/map2/www.birding.hu>

Breeding long-term trend: Barta, Z.: Vízirígó In: Haraszthy, L. (szerk.) (1998): Magyarország madarai. 441 p. National park directorates' databases (Annual survey of colonially breeding and strictly protected bird species) <http://map.mme.hu/maps/map2/www.birding.hu>

Republic of Ireland

Breeding population size: Balmer, D., Gillings, S., Caffrey, B., Swan, B., Downie, I. & Fuller, R. (2013) Bird Atlas 2007-11 The breeding and wintering birds of Britain and Ireland. British Trust for Ornithology. Gibbons D.W., Reid J.B. & Chapman R.A. (1993) The New Atlas of Breeding Birds in Britain and Ireland 1988-1991. Poyser, London.

Breeding short-term trend: Balmer, D., Gillings, S., Caffrey, B., Swan, B., Downie, I. & Fuller, R. (2013) Bird Atlas 2007-11 The breeding and wintering birds of Britain and Ireland. British Trust for Ornithology. Gibbons D.W., Reid J.B. & Chapman R.A. (1993) The New Atlas of Breeding Birds in Britain and Ireland 1988-1991. Poyser, London.

Breeding long-term trend: Balmer, D., Gillings, S., Caffrey, B., Swan, B., Downie, I. & Fuller, R. (2013) Bird Atlas 2007-11 The breeding and wintering birds of Britain and Ireland. British Trust for Ornithology. Sharrock, J.T.R. (1976) The Atlas of Breeding Birds in Britain and Ireland. T. & AD Poyser.

Italy

Breeding population size: Brichetti P & Fracasso G. 2007. Ornitologia italiana. Vol.4 (Apodidae-Prunellidae). Alberto Perdisa Editore, Bologna

Breeding short-term trend: No recent data available

Breeding long-term trend: Brichetti P., Meschini E., 1993. Stima delle popolazioni di uccelli nidificanti. In Meschini E., Frugis S., 1993. Atlante degli uccelli nidificanti in Italia. Suppl. Ric. Biol. Selvaggina, 20, 1-345.

Kosovo

Breeding population size: Qenan Maxhuni

Breeding short-term trend: Qenan Maxhuni

Breeding long-term trend: Puzovic, S. et al. (2004): Birds of Serbia and Montenegro – Size of nesting populations. I trends: 1990-2002. Ciconia 12

Latvia

Breeding population size: Unpublished data for European Breeding Bird Atlas (2013-2017); Expert: Andris Dekants, andris.dekants@lob.lv Expert: Aigars Kalvans, a.kalvans@lvm.lv

Breeding short-term trend: Unpublished data for European Breeding Bird Atlas (2013-2017); Expert: Andris Dekants, andris.dekants@lob.lv

Breeding long-term trend: Strazds M., Priednieks J., Vaverins G. 1994. [Size of Latvian bird populations.] (in Latvian) In: Putni dabā, 4: 3–18 Unpublished data for European Breeding Bird Atlas (2013-2017); Expert: Andris Dekants, andris.dekants@lob.lv

Luxembourg

Breeding population size: Hourlay F. (2009): Recensements des Cincles plongeurs *Cinclus cinclus* et des autres oiseaux d'eau présents sur l'Attert (Avril 2005). *Regulus Wissenschaftliche Berichte*, 24: 55-60; Konter A. (2010): Wasservogel an der Sauer im Raum Echternach: Bestand und Ausblick. *Wissenschaftliche Berichte*, 25: 41-55; Ornitho.lu (2018): online database *natur&environnement asbl* & Dachverband Deutscher Avifaunisten (DDA) e.V.; Luxembourg Recorder (2018): database Musée national d'histoire naturelle; Luxembourg Lorgé P., E. Melchior (2016): Die Vögel Luxemburgs. *Natur&environnement Luxembourg*. ISBN: 978-2-919920-01-3

Breeding short-term trend: Hourlay F. (2009): Recensements des Cincles plongeurs *Cinclus cinclus* et des autres oiseaux d'eau présents sur l'Attert (Avril 2005). *Regulus Wissenschaftliche Berichte*, 24: 55-60; Konter A. (2010): Wasservogel an der Sauer im Raum Echternach: Bestand und Ausblick. *Wissenschaftliche Berichte*, 25: 41-55; Ornitho.lu (2018): online database *natur&environnement asbl* & Dachverband Deutscher Avifaunisten (DDA) e.V.; Luxembourg Recorder (2018): database Musée national d'histoire naturelle; Luxembourg Lorgé P., E. Melchior (2016): Die Vögel Luxemburgs. *Natur&environnement Luxembourg*. ISBN: 978-2-919920-01-3; LUXOR (2018): *natur&environnement – Bird-database*, Luxembourg

Cinclus cinclus (White-throated Dipper)

Luxembourg

Breeding long-term trend: Hourlay F. (2009): Recensements des Cincles plongeurs *Cinclus cinclus* et des autres oiesaux d'eau présents sur l'Attert (Avril 2005). *Regulus Wissenschaftliche Berichte*, 24: 55-60; Konter A. (2010): Wasservogel an der Sauer im Raum Echternach: Bestand und Ausblick. *Wissenschaftliche Berichte*, 25: 41-55; Ornitho.lu (2018): online database natur&environnement asbl & Dachverband Deutscher Avifaunisten (DDA) e.V.; Luxembourg Recorder (2018): database Musée national d'histoire naturelle; Luxembourg Lorgé P., E. Melchior (2016): Die Vögel Luxemburgs. *Natur&environment Luxembourg*. ISBN: 978-2-919920-01-3; LUXOR (2018): natur&environment – Bird-database, Luxembourg

North Macedonia

Breeding population size: unpublished data from the European Breeding Bird Atlas 2

Breeding short-term trend: unpublished data from the European Breeding Bird Atlas 2

Montenegro

Breeding population size: Puzović, S., Simić, D., Saveljić, D., Gergelj, J., Tucakov, M., Stojnić, N., Hulo, I., Ham, I., Vizi, O., Sciban, M., Ruzić, M., Vucanović, M., Jovanović, T. (2004): Birds of Serbia and Montenegro – Size of nesting populations. *I trends: 1990-2002*. *Ciconia* 12,

Norway

Breeding population size: Shimmings P. & Øien, I.J. 2015. Bestandsestimater og trender for norske hekkefugler. NOF-rapport 2015-2.

Breeding long-term trend: Shimmings, P. & Øien, I.J. 2015. Bestandsestimater for norske hekkefugler. NOF Rapport 2-2015. 268 pp.

Poland

Breeding population size: Chodkiewicz T., Kuczyński L., Sikora A., Chylarecki P., Neubauer G., Ławicki Ł., Stawarczyk T. 2015. Ocena liczebności populacji ptaków lęgowych w Polsce w latach 2008–2012. *Ornis Polonica* 56: 149-189; Wilk T., Bobrek R., Pępkowska-Król A., Neubauer G., Kosicki J. 2016. Ptaki polskich Karpat - stan, zagrożenia, ochrona. OTOP, Marki.

Breeding short-term trend: Chief Inspectorate of Environmental Protection & Polish Society for the Protection of Birds (OTOP) / BirdLife Poland

Breeding long-term trend: Chief Inspectorate of Environmental Protection & Polish Society for the Protection of Birds (OTOP) / BirdLife Poland

Portugal

Breeding population size: eBird (2019). eBird: An online database of bird distribution and abundance [web application]. eBird, Ithaca, New York. Available: <http://www.ebird.org/portugal/home>. (Accessed: October 22, 2018).

Breeding short-term trend: eBird: An online database of bird distribution and abundance [web application]. eBird, Ithaca, New York. Available: <http://www.ebird.org/po>

Romania

Breeding population size: Ornitodata (Romanian Ornithological Society) Database, OpenBirdMaps (Milvus Group) Database, Rombird (Romanian Rarity Commission) Database

Breeding short-term trend: Ornitodata (Romanian Ornithological Society) Database, OpenBirdMaps (Milvus Group) Database, Rombird (Romanian Rarity Commission) Database

Breeding long-term trend: Ornitodata (Romanian Ornithological Society) Database, OpenBirdMaps (Milvus Group) Database, Rombird (Romanian Rarity Commission) Database

Russia

Breeding population size: Voltzit & Kalyakin 2013-2019; Database of the project on Atlas of breeding birds of European Russia

Breeding short-term trend: Belik unpublished. vpbelik@mail.ru

Breeding long-term trend: Belik et al. 2003

Serbia

Breeding population size: EBBA2 project; Puzović, S., Radišić, D., Ružić, M., Rajković, D., Radaković, M., Pantović, U., Janković, M., Stojnić, N., Šćiban, M., Tucakov, M., Gergelj, J., Sekulić, G., Agošton, A. & Raković, M. 2015. Birds of Serbia: Breeding Population Estimates and Trends for the Period 2008-2013. Bird protection and study society of Serbia, and Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad, Novi Sad.

Breeding short-term trend: EBBA2 project; Puzović, S., Radišić, D., Ružić, M., Rajković, D., Radaković, M., Pantović, U., Janković, M., Stojnić, N., Šćiban, M., Tucakov, M., Gergelj, J., Sekulić, G., Agošton, A. & Raković, M. 2015. Birds of Serbia: Breeding Population Estimates and Trends for the Period 2008-2013. Bird protection and study society of Serbia, and Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad, Novi Sad.

Breeding long-term trend: Puzović, S., Radišić, D., Ružić, M., Rajković, D., Radaković, M., Pantović, U., Janković, M., Stojnić, N., Šćiban, M., Tucakov, M., Gergelj, J., Sekulić, G., Agošton, A. & Raković, M. 2015. Birds of Serbia: Breeding Population Estimates and Trends for the Period 2008-2013. Bird protection and study society of Serbia, and Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad, Novi Sad.

Winter population size: IWC database; Šćiban, M., Grubač, B., Vujić, A. eds. 2018. Red book of fauna of Serbia III - Birds. Belgrade: Institute for Nature Conservation of Serbia, University of Novi Sad, Faculty of Sciences, Department of Biology and Ecology and Bird Protection and Study Society of Serbia.

Winter short-term trend: IWC database; Radišić, D., Vasić, V., Puzović, S., Ružić, M., Šćiban, M., Grubač, B., Vujić, A. eds. 2018. Red book of fauna of Serbia III - Birds. Belgrade: Institute for Nature Conservation of Serbia, University of Novi Sad, Faculty of Sciences, Department of Biology and Ecology and Bird Protection and Study Society of Serbia.

Winter long-term trend: IWC database; Bioras database <http://www.bioras.petnica.rs/home.php>

Slovakia

Breeding population size: Coordinatory group for reporting 2019. Danko Štefan, Darolová Alžbeta, Krištín Anton: Rozšírenie vtákov na Slovensku. VEDA, vyd. SAV Bratislava, 2002.

Breeding short-term trend: Coordinatory group for reporting 2019, AVES-Symfony Database 2013-2018, KIMS Database 2013-2018. Danko Štefan, Darolová Alžbeta, Krištín Anton: Rozšírenie vtákov na Slovensku. VEDA, vyd. SAV Bratislava, 2002.

Cinclus cinclus (White-throated Dipper)

Slovakia

Breeding long-term trend: Coordinatory group for reporting 2019, AVES-Symfony Database 2013-2018, KIMS Database 2013-2018. Danko Štefan, Darolová Alžbeta, Krištín Anton: Rozšírenie vtákov na Slovensku. VEDA, vyd. SAV Bratislava, 2002.

Slovenia

Breeding population size: MIHELIČ T., KMECL P., DENAC K., KOCE U., VREZEC A., DENAC D. (eds.) (2019): Atlas ptic Slovenije. Popis gnezdičk 2002–2017. – DOPPS, Ljubljana.

Breeding short-term trend: There are no sources for this information.

Breeding long-term trend: There are no sources for this information.

Spain

Breeding population size: Información proporcionada por las comunidades autónomas. Martí, R. & del Moral, J.C. (Eds.) (2003). Atlas de las Aves Reproductoras de España. Dirección General de Conservación de la Naturaleza- Sociedad Española de Ornitología. Madrid, 733 pp. (https://www.miteco.gob.es/es/biodiversidad/temas/inventarios-nacionales/inventario-especies-terrestres/inventario-nacional-de-biodiversidad/ieet_aves_atlas.aspx) Moreno-Rueda, G. (2016). Mirlo acuático – *Cinclus cinclus*. En: Enciclopedia Virtual de los Vertebrados Españoles. Salvador, A. & Morales, M.B. (Eds.). Museo Nacional de Ciencias Naturales, Madrid. (<http://www.vertebradosibericos.org/>)

Breeding short-term trend: Database of the 'Atlas de las aves reproductoras de España'. Updated version 2011 with data from SEO/Birdlife's monitoring programmes. In: Inventario Español de Especies Terrestres, Inventario Español del Patrimonio Natural y de la Biodiversidad. Ministerio de Agricultura, Alimentación y Medio Ambiente (2013). (https://www.miteco.gob.es/fr/biodiversidad/temas/inventarios-nacionales/inventario-especies-terrestres/ieet_aves_sist_seg_tendencia_comunes_esp.aspx) Información proporcionada por las Comunidades Autónomas. Moreno-Rueda, G. (2016). Mirlo acuático – *Cinclus cinclus*. En: Enciclopedia Virtual de los Vertebrados Españoles. Salvador, A. & Morales, M.B. (Eds.). Museo Nacional de Ciencias Naturales, Madrid. (<http://www.vertebradosibericos.org/>) SEO/BirdLife (2018). Programas de seguimiento y grupos de trabajo de SEO/Birdlife 2017. SEO/Birdlife. Madrid. 69 pp. (<https://www.seo.org/boletin/seguimiento/boletin/2017/html5forpc.html?page=0>). SEO/BirdLife (2019). Programas de seguimiento y grupos de trabajo de SEO/BirdLife 2018. SEO/BirdLife. Madrid. (<https://doi.org/10.31170/0073>)

Breeding long-term trend: Información proporcionada por las Comunidades Autónomas. Martí, R. & del Moral, J.C. (Eds.) (2003). Atlas de las Aves Reproductoras de España. Dirección General de Conservación de la Naturaleza- Sociedad Española de Ornitología. Madrid, 733 pp. (https://www.miteco.gob.es/es/biodiversidad/temas/inventarios-nacionales/inventario-especies-terrestres/inventario-nacional-de-biodiversidad/ieet_aves_atlas.aspx) Moreno-Rueda, G. (2016). Mirlo acuático – *Cinclus cinclus*. En: Enciclopedia Virtual de los Vertebrados Españoles. Salvador, A. & Morales, M.B. (Eds.). Museo Nacional de Ciencias Naturales, Madrid. (<http://www.vertebradosibericos.org/>)

Sweden

Breeding population size: Ottosson, U., Ottvall, R., Elmberg, J., Green, M., Gustafsson, R., Haas, F., Holmqvist, N., Lindström, Å., Nilsson, L., Svensson, M., Svensson, S. & Tjernberg, M. 2012. Fåglarna i Sverige – antal och förekomst. SOF, Halmstad. Swedish Bird Survey. BirdLife Sverige, Annual Bird reports.

Breeding short-term trend: Svensk fågeltaxering - Swedish Bird Survey

Breeding long-term trend: Svensk fågeltaxering - Swedish Bird Survey

Switzerland

Breeding population size: Knaus, P., S. Antoniazza, S. Wechsler, J. Guélat, M. Kéry, N. Strelbel & T. Sattler (2018): Swiss Breeding Bird Atlas 2013–2016. Distribution and population trends of birds in Switzerland and Liechtenstein. Swiss Ornithological Institute, Sempach.

Breeding short-term trend: <https://www.vogelwarte.ch/en/projects/population-trends/breeding-population-indices/>

Breeding long-term trend: <https://www.vogelwarte.ch/en/projects/population-trends/breeding-population-indices/>

Turkey

Breeding population size: Levent Erkol, , Ferdi Akarsu, Güven Eken personal communication (2019), Kusbank Bird Database (Ebird) Kirwan G.M., Boyla K. A., Castell P., Demirci B., Özen M., Welch H., Marlow T., 2008, Birds of Turkey. Londra, Christopher Helm, 978-1-4081-0475-

Ukraine

Breeding population size: Atlas work, non-published data

United Kingdom

Breeding population size: Baseline = Gibbons, D.W., Reid, J.B. & Chapman, R.A. 1993. The New Atlas of Breeding Birds in Britain and Ireland: 1988-1991. Poyser, London. Extrapolation from 1988-91 using Breeding Bird Survey monitoring trend.

Breeding short-term trend: BTO Waterways Bird Survey/Waterways Breeding Bird Survey <https://www.bto.org/about-birds/birdtrends/2018>

Breeding long-term trend: BTO Waterways Bird Survey/Waterways Breeding Bird Survey <https://www.bto.org/about-birds/birdtrends/2018>

Bibliography

- Bird, J. P., Martin, R., Akçakaya, H. R., Gilroy, J., Burfield, I. J., Garnett, S. G., Symes, A., Taylor, J., Sekercioglu, Ç. H. and Butchart, S. H. M. 2020. Generation lengths of the world's birds and their implications for extinction risk. *Conservation Biology* 34(5): 1252-1261. DOI: 10.1111/cobi.13486.
- BirdLife International. 2004. *Birds in Europe: population estimates, trends and conservation status*. BirdLife International, Cambridge, U.K.
- Crick, H.Q.P. and Sparks, T.H. 1999. Climate change related to egg-laying trends. *Nature* 399: 423-424.
- Crick, H.Q.P., Dudley, C., Glue, D.E. and Thomson, D.L. 1997. UK birds are laying earlier. *Nature* 388: 526.
- Flint, P.R. and Stewart, P.F. 1983. *The birds of Cyprus*. British Ornithologists' Union, London.
- Ormerod, S., Tyler, S. and Christie, D.A. 2015. White-throated Dipper (*Cinclus cinclus*). In: J. del Hoyo, A. Elliott, J. Sargatal, D.A. Christie & E. de Juana (eds), *Handbook of the Birds of the World Alive*, Lynx Edicions, Barcelona.
- Taylor, A.J. and O'Halloran, J. 2001. Diet of Dippers *Cinclus cinclus* during an early winter spate and the possible implications for Dipper populations subjected to climate change. *Bird Study* 48: 173-179.
- del Hoyo, J., Elliott, A. and Christie, D. 2005. *Handbook of the Birds of the World, Vol. 10: Cuckoo-shrikes to Thrushes*. Lynx Edicions, Barcelona, Spain.