Phoenicurus erythrogastrus (White-winged Redstart)

European Red List of Birds

Supplementary Material

The European Union (EU28) Red List assessments were based principally on the official data reported by EU Member States to the European Commission under Article 12 of the Birds Directive in 2019-20. For the European Red List assessments, similar data were sourced from BirdLife Partners and other collaborating experts in other European countries and territories. For more information, see BirdLife International (2021).

Contents

Reported national population sizes and trends
Trend maps of reported national population data
Sources of reported national population data
Species factsheet bibliography

Recommended citation

BirdLife International (2021) European Red List of Birds. Luxembourg: Publications Office of the European Union.

Further information

http://datazone.birdlife.org/info/euroredlist
http://www.birdlife.org/europe-and-central-asia/european-red-list-birds-0
http://www.iucnredlist.org/regions/europe
http://ec.europa.eu/environment/nature/conservation/species/redlist/

Data requests and feedback

To request access to these data in electronic format, provide new information, correct any errors or provide feedback, please email science@birdlife.org.

Phoenicurus erythrogastrus (White-winged Redstart)

Table 1. Reported national breeding population size and trends in Europe[1].

Country (or territory)[2]	Population estimate				Short-term population trend[5]				Long-term population trend[5]				Subspecific population (where relevant)
	Size (pairs)[3]	Europe (%)	Year(s)	Method[4]	Direction[6]	Magnitude (%)[7]	Year(s)	Method[4]	Direction[6]	Magnitude (%)[7]	Year(s)	Method[4]	
Armenia	0	<1	2013-2018	complete	?		2007-2018		?		2003-2018	deficient	
Georgia	530–5300	51	2013-2017	partial	?			deficient	?				
Russia	1200–2200	49	2008-2018	partial	-	-49 to -30	2008-2018	expert	-	-49 to -30	1980-2018	expert	
Europe	**1700–7500**	**100**											

[1] See 'Sources' at end of factsheet, and for more details on individual EU Member State reports, see the Article 12 reporting portal at http://bd.eionet.europa.eu/article12/report.
[2] The designation of geographical entities and the presentation of the material do not imply the expression of any opinion whatsoever on the part of IUCN or BirdLife International concerning the legal status of any country, territory or area, or of its authorities, or concerning the delimitation of its frontiers or boundaries.
[3] In the few cases where population size estimates were reported in units other than those specified, they were converted to the correct units using standard correction factors.
[4] The 'method used' (replacing the data 'quality' assessment in the 2015 European Red List) is reported as: a) Complete: complete survey or a statistically robust estimate; b) Partial: based mainly on extrapolation from a limited amount of data; c) Expert: based mainly on expert opinion with very limited data; d) Defficient: insufficient or no data available.
[5] The robustness of regional trends to the effects of any missing or incomplete data was tested using plausible scenarios, based on other sources of information, including any other reported information, recent national Red Lists, scientific literature, other publications and consultation with relevant experts.
[6] Trend directions are reported as: increasing (+); decreasing (-); stable (0); fluctuating (F); or unknown (?).
[7] Trend magnitudes are rounded to the nearest integer.

Trend maps

A symbol appears in each country where the species occurs: the shape and colour of the symbol represent the population trend in that country, and the size of the symbol corresponds to the proportion of the European population occurring in that country.

KEY

- Large increase (≥50%)
- Moderate increase (20–49%)
- Small increase (<20%)
- Increase of unknown magnitude
- Large decrease (≥50%)
- Moderate decrease (20–49%)
- Small decrease (<20%)
- Decrease of unknown magnitude
- Stable or fluctuating
- Unknown
- Present (no population or trend data)
- Extinct since 1980

Each symbol, with the exception of Present and Extinct, may occur in up to three different size classes, corresponding to the proportion of the European population occurring in that country.

- Large: ≥10% of the European population
- Medium: 1–9% of the European population
- Small: <1% of the European population

The designation of geographical entities and the presentation of the material do not imply the expression of any opinion whatsoever on the part of IUCN or BirdLife International concerning the legal status of any country, territory or area, or of its authorities, or concerning the delimitation of its frontiers or boundaries.

Figure 1. Breeding population sizes and short-term trends across Europe.

Figure 2. Breeding population sizes and long-term trends across Europe.

Sources

Armenia

Breeding population size: TSE NGO National Bird Monitoring data.
Breeding short-term trend: TSE (2020) The Atlas of the Breeding Birds in Armenia. In preparation.
Breeding long-term trend: TSE (2020) The Atlas of the Breeding Birds in Armenia. In preparation.

Georgia

Breeding population size: EBBA Georgia, prepared by Sabuko-Society for nature conservation, Ilia state university and NGO "Psovi"

Russia

Breeding population size: Voltzit & Kalyakin 2013-2019; Database of the project on Atlas of breeding birds of European Russia
Breeding short-term trend: Belik unpublished
Breeding long-term trend: Belik et al. 2003

Bibliography

Bird, J. P., Martin, R., Akçakaya, H. R., Gilroy, J., Burfield, I. J., Garnett, S. G., Symes, A., Taylor, J., Sekercioglu, Ç. H. and Butchart, S. H. M. 2020. Generation lengths of the world's birds and their implications for extinction risk. Conservation Biology 34(5): 1252-1261. DOI: 10.1111/cobi.13486.

BirdLife International. 2004. Birds in Europe: population estimates, trends and conservation status. BirdLife International, Cambridge, U.K.

Brazil, M. 2009. Birds of East Asia: eastern China, Taiwan, Korea, Japan, eastern Russia. Christopher Helm, London.

Collar, N. 2015. Güldenstädt's Redstart (Phoenicurus erythrogastrus). In: del Hoyo, J., Elliott, A., Sargatal, J., Christie, D.A. and de Juana, E. (eds), Handbook of the Birds of the World Alive, Lynx Edicions, Barcelona.

Heath, M. and Tucker, G. 1994. Birds in Europe. World Birdwatch 16: 9-13.